# IASI Principle Component Analysis, IASI and AIRS Regression Retrieval Comparisons and Update on Radiance Climatology Dataset


Mitchell D. Goldberg
Zhaohui Cheng
Lihang Zhou
Chris Barnet

NOAA/NESDIS AIRS Science Team October 9, 2007


#### **IASI PCA** Motivation


- IASI PCA Regression
- Data Compression
- Noise Estimation
- Detection of Anomalies
- Assimilation of Reconstructed Radiances
- Assimilation of PCs

## Spectral Coverage of Thermal Sounders (Example BT's for AIRS, IASI, & CrIS)




## Instrument Noise, NEΔT at 250 K (Interferometers Noise Is Apodized)








#### Generation of EOFs


- 3 bands together
- Separate Bands
  - 645-1210 cm-1
  - -1210 2000
  - **2000 2760**


































IASI [670.000cm-1] Granule-125 20070910

IASI [670.000cm-1] Granule-125 20070910




IASI [770.000cm-1] Granule-125 20070910

IASI [770.000cm-1] Granule-125 20070910




IASI [900.000cm-1] Granule-125 20070910

IASI [900.000cm-1] Granule-125 20070910





IASI [965.000cm-1] Granule-125 20070910

IASI [965.000cm-1] Granule-125 20070910





IASI [1041.50cm-1] Granule-125 20070910






IASI [1298.00cm-1] Granule-125 20070910







IASI [1298.00cm-1] Granule-125 20070910



IASI [1306.25cm-1] Granule-125 20070910



IASI [1594.75cm-1] Granule-125 20070910





1.06

-5.36


-3.75


-2.15

-2


٥


D.01







IASI [1739.00cm-1] Granule-125 20070910







IASI [1923.00cm-1] Granule-125 20070910







IASI [1923.00cm-1] Granule-125 20070910







260 Rec BT (K) 220 240 IASI [2141cm-1] Granule-125 20070910






IASI [2141cm-1] Granule-125 20070910






IASI [2375cm-1] Granule-125 20070910

IASI [2375cm-1] Granule-125 20070910




### IASI and AIRS Regression Temperature Retrieval Accuracy



Independent dataset (ocean only)


# IASI and AIRS Regression Moisture Retrieval Accuracy




Independent dataset (ocean only)

### IASI


### **AIRS**



### AIRS Regression Retrieval— Temperature Profile (opr coef vs exp)




### AIRS Regression Retrieval— Temperature Profile (opr vs exp)



Independent dataset (ocean only)

Time Series of Ret. Temp.(0.76 mb) Avg. of -90 to 90 266 266 264 282 260 258 255 254 252 250 OCT JÁN APR ASEMBOLAROS (Final) DESCENDING (Final) OCT JÁN APR JÚL ASCÉMENG (ECMWF) DESCENDING (ECMWF) JÁN APR 2007ASCENDING (Res DESCENDING (Re JÚL ост ост


Time Series of Ret. Temp.(5 mb)
Avg. of -90 to 90





# **Generation of AIRS Radiance Dataset for Climate Studies**

# Applications of Mapped Spectrally Resolved Radiances

- Compare radiances with simulated radiances from model analyses
- Compare different years to see how the outgoing infrared radiances have changed.









#### Water – Upper Troposphere

1520.87cm-1





Radiance Climate Dataset detected jump in ECMWF moisture analysis after update to their adaptive bias tuning


#### Observed AIRS minus ECMWF Simulated AIRS for Upper Trop. Water Vapor




#### Observed AIRS minus NCEP Simulated AIRS for Upper Trop. Water Vapor










### Consistent with Sept 03,04,05








Limb Adjusted BT, 7 PCs - ECHWF (NAD), 1519.07cm-1, Clear Sky, Jul, 2008






27007-08-17


NOAA/NESDIS/STAR/SMCD/SPB/IOSSPDT

2007-08-17 19:07


## GDAS consistent with Sept 05


Limb Adjusted BT, 7 PCs - GDAS (NAD), 1519.07cm-1, Clear Sky, Jan, 2006





Limb Adjusted BT, 7 PCs - GDAS (NAD), 1519.07cm-1, Clear Sky, Jul, 2006







NOAA/NESDIS/STAR/SMCD/SPB/IOSSPDT


### Consistent with Sept 03,04,05


Limb Adjusted BT, 7 PCs - ECMWF (NAD), 1598, 9cm-1, Clear Sky, Jan, 2006





Limb Adjusted BT, 7 PCs - ECMWF (NAD), 1598.49cm-1, Clear Sky, Jul, 2008





# Summary

- Continue IASI PCA research
- Compare IASI and AIRS retrievals
- Complete 2003-2007 AIRS radiance climate dataset.
- Begin generating IASI radiance climate dataset.
- IASI level 2 product will be operational at NESDIS in June 2008

## Summary

- IASI and AIRS are critical for satellite intercalibration/characterization of other key observing systems (e.g International Geostationary satellites)
- New WMO Global Space-based Intercalibration System (GSICS) led by NESDIS
- Establish stability of IASI using AIRS

## Organizations contributing to GSICS

- NOAA
- NIST
- NASA
- EUMETSAT
- CNES
- CMA
- JMA
- KMA

GSICS current focus is on the intercalibration of operational satellites, and makes use of key research instruments such as AIRS and MODIS to intercalibration the operational instruments

